
https://answersun.com/download/
solutions-manual-of-calculus-one-and-several-variables-by-salas-saturnino-10th-edition/

TENTH
EDITION



SALAS
HILLE
ETGEN

CALCULUS


ONE VARIABLE

1807 WILEY 2007

JOHN WILEY & SONS, INC.

Download full file from answersun.com


https://answersun.com/download/
solutions-manual-of-calculus-one-and-several-variables-by-salas-saturnino-10th-edition/



CONTENTS


CHAPTER 1 PRECALCULUS REVIEW 1

1.1 What is Calculus? 1
1.2 Review of Elementary Mathematics 3
1.3 Review of Inequalities 11
1.4 Coordinate Plane; Analytic Geometry 17
1.5 Functions 24
1.6 The Elementary Functions 32
1.7 Combinations of Functions 41
1.8 A Note on Mathematical Proof; Mathematical Induction 47




CHAPTER 2 LIMITS AND CONTINUITY 53

2.1 The Limit Process (An Intuitive Introduction) 53
2.2 Definition of Limit 64
2.3 Some Limit Theorems 73
2.4 Continuity 82
2.5 The Pinching Theorem; Trigonometric Limits 91
2.6 Two Basic Theorems 97
Project 2.6 The Bisection Method for Finding the Roots of $f(x) = 0$ 102




Download full file from answersun.com

https://answersun.com/download/
solutions-manual-of-calculus-one-and-several-variables-by-salas-saturnino-10th-edition/




CHAPTER 3 THE DERIVATIVE; THE PROCESS OF DIFFERENTIATION 105

3.1 The Derivative 105
3.2 Some Differentiation Formulas 115
3.3 The *d/dx* Notation; Derivatives of Higher Order 124
3.4 The Derivative As A Rate of Change 130
3.5 The Chain Rule 133
3.6 Differentiating The Trigonometric Functions 142
3.7 Implicit Differentiation; Rational Powers 147



CHAPTER 4 THE MEAN-VALUE THEOREM; APPLICATIONS OF THE FIRST AND SECOND DERIVATIVES 154

4.1 The Mean-Value Theorem 154
4.2 Increasing and Decreasing Functions 160
4.3 Local Extreme Values 167
4.4 Endpoint Extreme Values; Absolute Extreme Values 174
4.5 Some Max-Min Problems 182
Project 4.5 Flight Paths of Birds 190
4.6 Concavity and Points of Inflection 190
4.7 Vertical and Horizontal Asymptotes; Vertical Tangents and Cusps 195
4.8 Some Curve Sketching 201
4.9 Velocity and Acceleration; Speed 209
Project 4.9A Angular Velocity; Uniform Circular Motion 217
Project 4.9B Energy of a Falling Body (Near the Surface of the Earth) 217
4.10 Related Rates of Change per Unit Time 218
4.11 Differentials 223
Project 4.11 Marginal Cost, Marginal Revenue, Marginal Profit 228
4.12 Newton-Raphson Approximations 229



CHAPTER 5 INTEGRATION 234


5.1 An Area Problem; a Speed-Distance Problem 234
5.2 The Definite Integral of a Continuous Function 237
5.3 The Function $f(x) = \int_a^x f(t)\,dt$ 246
5.4 The Fundamental Theorem of Integral Calculus 254
5.5 Some Area Problems 260
Project 5.5 Integrability; Integrating Discontinuous Functions 266
5.6 Indefinite Integrals 268
5.7 Working Back from the Chain Rule; the *u*-Substitution 274
5.8 Additional Properties of the Definite Integral 281
5.9 Mean-Value Theorems for Integrals; Average Value of a Function 285

Download full file from answersun.com

https://answersun.com/download/
solutions-manual-of-calculus-one-and-several-variables-by-salas-saturnino-10th-edition/


CHAPTER 6 SOME APPLICATIONS OF THE INTEGRAL 292

6.1 More on Area 292
6.2 Volume by Parallel Cross Sections; Disks and Washers 296
6.3 Volume by the Shell Method 306
6.4 The Centroid of a Region; Pappus's Theorem on Volumes 312
Project 6.4 Centroid of a Solid of Revolution 319
6.5 The Notion of Work 319
*6.6 Fluid Force 327




CHAPTER 7 THE TRANSCENDENTAL FUNCTIONS 333

7.1 One-to-One Functions; Inverses 333
7.2 The Logarithm Function, Part I 342
7.3 The Logarithm Function, Part II 347
7.4 The Exponential Function 356
Project 7.4 Some Rational Bounds for the Number e 364
7.5 Arbitrary Powers; Other Bases 364
7.6 Exponential Growth and Decay 370
7.7 The Inverse Trigonometric Functions 378
Project 7.7 Refraction 387
7.8 The Hyperbolic Sine and Cosine 388
*7.9 The Other Hyperbolic Functions 392




CHAPTER 8 TECHNIQUES OF INTEGRATION 398

8.1 Integral Tables and Review 398
8.2 Integration by Parts 402
Project 8.2 Sine Waves $y = \sin nx$ and Cosine Waves $y = \cos nx$ 410
8.3 Powers and Products of Trigonometric Functions 411
8.4 Integrals Featuring $\sqrt{a^2 - x^2}$, $\sqrt{a^2 + x^2}$, $\sqrt{x^2 - a^2}$ 417
8.5 Rational Functions; Partial Fractions 422
*8.6 Some Rationalizing Substitutions 430
8.7 Numerical Integration 433



CHAPTER 9 SOME DIFFERENTIAL EQUATIONS 443


9.1 First-Order Linear Equations 444
9.2 Integral Curves; Separable Equations 451
Project 9.2 Orthogonal Trajectories 458
9.3 The Equation $y'' + ay' + by = 0$ 459



*Denotes optional section.

Download full file from answersun.com

https://answersun.com/download/
solutions-manual-of-calculus-one-and-several-variables-by-salas-saturnino-10th-edition/




CHAPTER 10 THE CONIC SECTIONS; POLAR COORDINATES; PARAMETRIC EQUATIONS 469

10.1 Geometry of Parabola, Ellipse, Hyperbola 469
10.2 Polar Coordinates 478
10.3 Sketching Curves in Polar Coordinates 484
Project 10.3 Parabola, Ellipse, Hyperbola in Polar Coordinates 491
10.4 Area in Polar Coordinates 492
10.5 Curves Given Parametrically 496
Project 10.5 Parabolic Trajectories 503
10.6 Tangents to Curves Given Parametrically 503
10.7 Arc Length and Speed 509
10.8 The Area of A Surface of Revolution; The Centroid of a Curve; Pappus's Theorem on Surface Area 517
Project 10.8 The Cycloid 525



CHAPTER 11 SEQUENCES; INDETERMINATE FORMS; IMPROPER INTEGRALS 528

11.1 The Least Upper Bound Axiom 528
11.2 Sequences of Real Numbers 532
11.3 Limit of a Sequence 538
Project 11.3 Sequences and the Newton-Raphson Method 547
11.4 Some Important Limits 550
11.5 The Indeterminate Form (0/0) 554
11.6 The Indeterminate Form (∞/∞); Other Indeterminate Forms 560
11.7 Improper Integrals 565



CHAPTER 12 INFINITE SERIES 575

12.1 Sigma Notation 575
12.2 Infinite Series 577
12.3 The Integral Test; Basic Comparison, Limit Comparison 585
12.4 The Root Test; the Ratio Test 593
12.5 Absolute Convergence and Conditional Convergence; Alternating Series 597
12.6 Taylor Polynomials in x; Taylor Series in x 602
12.7 Taylor Polynomials and Taylor Series in $x - a$ 613
12.8 Power Series 616
12.9 Differentiation and Integration of Power Series 623
Project 12.9A The Binomial Series 633
Project 12.9B Estimating π 634

Download full file from answersun.com

https://answersun.com/download/
solutions-manual-of-calculus-one-and-several-variables-by-salas-saturno-10th-edition/

APPENDIX A SOME ADDITIONAL TOPICS A-1

A.1 Rotation of Axes; Eliminating the xy-Term A-1
A.2 Determinants A-3

APPENDIX B SOME ADDITIONAL PROOFS A-8

B.1 The Intermediate-Value Theorem A-8
B.2 Boundedness; Extreme-Value Theorem A-9
B.3 Inverses A-10
B.4 The Integrability of Continuous Functions A-11
B.5 The Integral as the Limit of Riemann Sums A-14

ANSWERS TO ODD-NUMBERED EXERCISES A-15
Index I-1
Table of Integrals Inside Covers

Download full file from answersun.com

https://answersun.com/download/
solutions-manual-of-calculus-one-and-several-variables-by-salas-saturnino-10th-edition/

THEOREM 2.4.4

If g is continuous at c and f is continuous at $g(c)$, then the composition $f \circ g$ is continuous at c.

The idea here is as follows: with g continuous at c, we know that

$$\text{for } x \text{ close to } c,\ g(x) \text{ is close to } g(c);$$

from the continuity of f at $g(c)$, we know that

$$\text{with } g(x) \text{ close to } g(c),\ f(g(x)) \text{ is close to } f(g(c)).$$

In summary,

$$\text{with } x \text{ close to } c,\ f(g(x)) \text{ is close to } f(g(c)).$$

The argument we just gave is too vague to be a proof. Here, in contrast, is a proof. We begin with $\epsilon > 0$. We must show that there exists a number $\delta > 0$ such that

$$\text{if} \quad |x - c| < \delta, \quad \text{then} \quad |f(g(x)) - f(g(c))| < \epsilon.$$

In the first place, we observe that, since f is continuous at $g(c)$, there does exist a number $\delta_1 > 0$ such that

$$(1) \qquad \text{if} \quad |t - g(c)| < \delta_1, \quad \text{then} \quad |f(t) - f(g(c))| < \epsilon.$$

With $\delta_1 > 0$, we know from the continuity of g at c that there exists a number $\delta > 0$ such that

$$(2) \qquad \text{if} \quad |x - c| < \delta, \quad \text{then} \quad |g(x) - g(c)| < \delta_1.$$

Combining (2) and (1), we have what we want: by (2),

$$\text{if} \quad |x - c| < \delta, \quad \text{then} \quad |g(x) - g(c)| < \delta_1$$

so that by (1)

$$|f(g(x)) - f(g(c))| < \epsilon.$$

This proof is illustrated in Figure 2.4.5. The numbers within δ of c are taken by g to within δ_1 of $g(c)$, and then by f to within ϵ of $f(g(c))$.




Figure 2.4.5

It's time to look at some examples.

Download full file from answersun.com

https://answersun.com/download/
solutions-manual-of-calculus-one-and-several-variables-by-salas-saturnino-10th-edition/

division proposes that the road be constructed by restoring a section of the old road from A to some point P and constructing a new road from P to B. Given that the cost of restoring the old road is \$2,000,000 per mile and the cost of a new road is \$4,000,000 per mile, how much of the old road should be restored so as to minimize the cost of the project.




Figure 4.5.5

SOLUTION Figure 4.5.5 shows the geometry of the problem. Notice that we have chosen a straight line joining P and B rather than some curved path. (The shortest connection between two points is provided by the straight-line path.) We let x be the amount of old road that will be restored. Then

$$\sqrt{9+(5-x)^2} = \sqrt{34-10x+x^2}$$

is the length of the new part. The total cost of constructing the two sections of road is

$$C(x) = 2\cdot 10^6 x + 4\cdot 10^6[34-10x+x^2]^{1/2}, \qquad 0 \le x \le 5.$$

We want to find the value of x that minimizes this function.

Differentiation gives

$$C'(x) = 2\cdot 10^6 + 4\cdot 10^6\left(\tfrac{1}{2}\right)[34-10x+x^2]^{-1/2}(2x-10)$$

$$= 2\cdot 10^6 + \frac{4\cdot 10^6(x-5)}{[34-10x+x^2]^{1/2}}, \qquad 0 < x < 5.$$

Setting $C'(x) = 0$, we find that

$$1 + \frac{2(x-5)}{[34-10x+x^2]^{1/2}} = 0$$

$$2(x-5) = -[34-10x+x^2]^{1/2}$$

$$4(x^2-10x+25) = 34-10x+x^2$$


$$3x^2 - 30x + 66 = 0$$

$$x^2 - 10x + 22 = 0.$$

By the general quadratic formula, we have

$$x = \frac{10 \pm \sqrt{100-4(22)}}{2} = 5 \pm \sqrt{3}.$$

The value $x = 5+\sqrt{3}$ is not in the domain of our function; the value we want is $x = 5-\sqrt{3}$. We analyze the sign of C':



Since the function is continuous on $[0, 5]$, it decreases on $[0, 5-\sqrt{3}]$ and increases on $[5-\sqrt{3}, 5]$. The number $x = 5-\sqrt{3} \cong 3.27$ gives the minimum value of C. The highway department will minimize its costs by restoring 3.27 miles of the old road. ❑

Example 5 (*The angle of incidence equals the angle of reflection.*) Figure 4.5.6 depicts light from point A reflected by a mirror to point B. Two angles have been marked: the *angle of incidence*, θ_i, and the *angle of reflection*, θ_r. Experiment shows




Figure 4.5.6

Download full file from answersun.com

https://answersun.com/download/solutions-manual-calculus-one-and-several-variables-by-salas-saturnino-10th-edition/

33. (a) Let f be continuous on $[-a, 0]$. Use a change of variable to show that

$$\int_{-a}^{0} f(x)\,dx = \int_{0}^{a} f(-x)\,dx.$$

(b) Let f be continuous on $[-a, a]$. Show that

$$\int_{-a}^{a} f(x)\,dx = \int_{0}^{a} [f(x) + f(-x)]\,dx.$$

34. Let f be a function continuous on $[-a, a]$. Prove the statement basing your argument on Exercise 33.

(a) $\displaystyle\int_{-a}^{a} f(x)\,dx = 0$ if f is odd.

(b) $\displaystyle\int_{-a}^{a} f(x)\,dx = 2\int_{0}^{a} f(x)\,dx$ if f is even.

Exercises 35–38. Evaluate using symmetry considerations.

35. $\displaystyle\int_{-\pi/4}^{\pi/4} (x + \sin 2x)\,dx.$

36. $\displaystyle\int_{-3}^{3} \frac{t^3}{1+t^2}\,dt.$

37. $\displaystyle\int_{-\pi/3}^{\pi/3} (1 + x^2 - \cos x)\,dx.$

38. $\displaystyle\int_{-\pi/4}^{\pi/4} (x^2 - 2x \,\vdots\, \sin x \,\vdots\, \cos 2x)\,dx.$

■ 5.9 MEAN-VALUE THEOREMS FOR INTEGRALS; AVERAGE VALUE OF A FUNCTION

We begin with a result that we asked you to prove earlier. (Exercise 33, Section 5.3.)

THEOREM 5.9.1 THE FIRST MEAN-VALUE THEOREM FOR INTEGRALS

If f is continuous on $[a, b]$, then there is at least one number c in (a, b) for which

$$\int_{a}^{b} f(x)\,dx = f(c)(b - a).$$

This number $f(c)$ is called *the average value* (or *mean value*) *of* f *on* $[a, b]$.

We now have the following identity:

(5.9.2) $$\int_{a}^{b} f(x)\,dx = (\textit{the average value of } f \textit{ on } [a, b]) \cdot (b - a).$$



Figure 5.9.1

This identity provides a powerful, intuitive way of viewing the definite integral. Think for a moment about area. If f is constant and positive on $[a, b]$, then Ω, the region below the graph, is a rectangle. Its area is given by the formula

$$\text{area of } \Omega = (\text{the constant value of } f \text{ on } [a, b]) \cdot (b - a). \quad \text{(Figure 5.9.1)}$$

If f is now allowed to vary continuously on $[a, b]$, then we have

$$\text{area of } \Omega = \int_{a}^{b} f(x)\,dx,$$

and the area formula reads

$$\text{area of } \Omega = (\textit{the average value of } f \text{ on } [a, b]) \cdot (b - a). \quad \text{(Figure 5.9.2)}$$

Download full file from answersun.com

https://answersun.com/download/
solutions-manual-of-calculus-one-and-several-variables-by-salas-saturnino-10th-edition/




Figure 10.3.1

In Figure 10.3.2 we have marked the values of θ where r is zero and the values of θ where r takes on an extreme value.




Figure 10.3.2

Reading from the figure we have the following: as θ increases from 0 to $\frac{1}{3}\pi$, r increases from -1 to 0; as θ increases from $\frac{1}{3}\pi$ to π, r increases from 0 to 3; as θ increases from π to $\frac{5}{3}\pi$, r decreases from 3 to 0; finally, as θ increases from $\frac{5}{3}\pi$ to 2π, r decreases from 0 to -1.

By applying this information step by step, we develop a sketch of the curve $r = 1 - 2\cos\theta$ in polar coordinates. (Figure 10.3.3.)





Figure 10.3.3

Download full file from answersun.com

https://answersun.com/download/solutions-manual-of-calculus-one-and-several-variables-by-salas-saturnino-10th-edition/

that remains after all of the "middle thirds" have been deleted is called the *Cantor middle third set*. Give some points that are in the Cantor set.

29. Start with a square that has sides four units long. Join the midpoints of the sides of the square to form a second square inside the first. Then join the midpoints of the sides of the second square to form a third square, and so on. See the figure. Find the sum of the areas of the squares.



30. (a) Show that if the series $\sum a_k$ converges and the series $\sum b_k$ diverges, then the series $\sum(a_k + b_k)$ diverges.

(b) Give examples to show that if $\sum a_k$ and $\sum b_k$ both diverge, then each of the series

$$\sum(a_k + b_k) \qquad \text{and} \qquad \sum(a_k - b_k)$$

may converge or may diverge.

31. Let $\sum_{k=0}^{\infty} a_k$ be a convergent series and let $R_n = \sum_{k=n+1}^{\infty} a_k$. Prove that $R_n \to 0$ as $n \to \infty$. Note that if s_n is the nth partial sum of the series, then $\sum_{k=0}^{\infty} a_k = s_n + R_n$; R_n is called the *remainder*.

32. (a) Prove that if $\sum_{k=0}^{\infty} a_k$ is a convergent series with all terms nonzero, then $\sum_{k=0}^{\infty} (1/a_k)$ diverges.

(b) Suppose that $a_k > 0$ for all k and $\sum_{k=0}^{\infty} a_k$ diverges. Show by example that $\sum_{k=0}^{\infty} (1/a_k)$ may converge and it may diverge.

33. Show that

$$\sum_{k=1}^{\infty} \ln\left(\frac{k+1}{k}\right) \quad \text{diverges}$$

although

$$\ln\left(\frac{k+1}{k}\right) \to 0.$$

34. Show that

$$\sum_{k=1}^{\infty} \left(\frac{k+1}{k}\right)^k \quad \text{diverges.}$$

35. (a) Assume that $d_k \to 0$ and show that

$$\sum_{k=1}^{\infty} (d_k - d_{k+1}) = d_1.$$

(b) Sum the following series:

$$\text{(i)} \sum_{k=1}^{\infty} \frac{\sqrt{k+1} - \sqrt{k}}{\sqrt{k(k+1)}}. \qquad \text{(ii)} \sum_{k=1}^{\infty} \frac{2k+1}{k^2(k+1)^2}.$$

36. Show that

$$\sum_{k=1}^{\infty} k x^{k-1} = \frac{1}{(1-x)^2} \qquad \text{for} \qquad |x| < 1.$$

HINT: Verify that s_n, the nth partial sum of the series, satisfies the identity

$$(1-x)^2 s_n = 1 - (n+1)x^n + n x^{n+1}.$$

(Exercises 37–40. *Speed of convergence*) Find the least integer N for which the nth partial sum of the series differs from the sum of the series by less than 0.0001.

37. $\sum_{k=0}^{\infty} \frac{1}{4^k}$.

38. $\sum_{k=0}^{\infty} (0.9)^k$.

39. $\sum_{k=1}^{\infty} \frac{1}{k(k+2)}$.

40. $\sum_{k=0}^{\infty} \left(\frac{2}{3}\right)^k$.

41. Start with the geometric series $\sum_{k=0}^{\infty} x^k$ with $|x| < 1$ and a positive number ϵ. Determine the least positive integer N for which $|L - s_N| < \epsilon$ given that the sum of the series is L and s_N is the Nth partial sum.

42. Prove that the series $\sum_{k=2}^{\infty} (a_{k+1} - a_k)$ converges iff the a_n tend to a finite limit.

■ 12.3 THE INTEGRAL TEST; BASIC COMPARISON, LIMIT COMPARISON

Here we begin our study of *series with nonnegative terms*: $a_k \geq 0$ for all k. For such series the sequence of partial sums is nondecreasing:

$$s_{n+1} = \sum_{k=0}^{n+1} a_k = a_{n+1} + \sum_{k=0}^{n} a_k \geq \sum_{k=0}^{n} a_k = s_n.$$

Download full file from answersun.com

https://answersun.com/download/

solutions-manual-of-calculus-one-and-several-variables-by-salas-saturnino-10th-edition/

11. $\frac{1}{x^2+2x+2}$ 13. $\frac{2}{x\sqrt{4x^4-1}}$ 15. $\frac{2x}{\sqrt{1-4x^2}} + \arcsin 2x$ 17. $\frac{2\arcsin x}{\sqrt{1-x^2}}$ 19. $\frac{x-(1+x^2)\arctan x}{x^2(1+x^2)}$

21. $\frac{1}{(1+4x^2)\sqrt{\arctan 2x}}$ 23. $\frac{1}{x[1+(\ln x)^2]}$ 25. $-\frac{r}{|r|\sqrt{1-r^2}}$ 27. $2x\,\text{arcsec}\left(\frac{1}{x}\right) - \frac{x^2}{\sqrt{1-x^2}}$

29. $\cos[\text{arcsec}(\ln x)] \cdot \frac{1}{x|\ln x|\sqrt{(\ln x)^2-1}}$ 31. $\sqrt{\frac{c-x}{c+x}}$

33. (a) x (b) $\sqrt{1-x^2}$ (c) $\frac{x}{\sqrt{1-x^2}}$ (d) $\frac{\sqrt{1-x^2}}{x}$ (e) $\frac{1}{\sqrt{1-x^2}}$ (f) $\frac{1}{x}$ 35. $\arcsin\left(\frac{x+b}{a}\right) + C$

39. $\frac{1}{4}\pi$ 41. $\frac{1}{4}\pi$ 43. $\frac{1}{20}\pi$ 45. $\frac{1}{24}\pi$ 47. $\frac{1}{3}\,\text{arcsec}\,4 - \frac{\pi}{9}$ 49. $\frac{1}{6}\pi$ 51. $\arctan 2 - \frac{1}{4}\pi \cong 0.322$

53. $\frac{1}{6}\arcsin x^2 + C$ 55. $\frac{1}{2}\arctan x^2 + C$ 57. $\frac{1}{3}\arctan\left(\frac{1}{3}\tan x\right) + C$ 59. $\frac{1}{2}(\arcsin x)^2 + C$ 61. $\arcsin(\ln x) + C$

63. $\frac{1}{3}\pi$ 65. $2\pi - \frac{4}{3}$ 67. $4\pi(\sqrt{2}-1)$

69. $\sqrt{s^2+sk}$ feet from the point where the line of the sign intersects the road.

71. (b) $\frac{1}{2}\pi a^2$; area of semicircle of radius a 75. $\frac{1}{\sqrt{1-x^2}}$ is not defined for $x \geq 1$.

77. estimate $\cong 0.523$, $\sin 0.523 \cong 0.499$ explanation: the integral = $\arcsin 0.5$; therefore $\sin(\text{integral}) = 0.5$

SECTION 7.8

1. $2x\cosh x^2$ 3. $\frac{a\sinh ax}{2\sqrt{\cosh ax}}$ 5. $\frac{1}{1-\cosh x}$ 7. $ab(\cosh bx - \sinh ax)$ 9. $\frac{a\cosh ax}{\sinh ax}$ 11. $2e^{2x}\cosh(e^{2x})$

13. $-e^{-x}\cosh 2x + 2e^{-x}\sinh 2x$ 15. $\tanh x$ 17. $(\sinh x)^x[\ln(\sinh x) + x\coth x]$ 27. absolute max -3

31. $A = 2$, $B = \frac{1}{3}$, $c = 3$ 33. $\frac{1}{a}\sinh ax + C$ 35. $\frac{1}{3a}\sinh^3 ax + C$ 37. $\frac{1}{a}\ln(\cosh ax) + C$ 39. $-\frac{1}{a\coth ax} + C$

41. $\frac{1}{2}(\sinh x\cosh x + x) + C$ 43. $2\cosh\sqrt{x} + C$ 45. $\sinh 1 \cong 1.175$ 47. [illegible] 49. π

51. $\pi[\ln 5 + \frac{1}{4}\sinh(4\ln 5)] \cong 250.492$ 53. (a) (0.69315, 1.25)
(b) $A \cong 0.38629$

SECTION 7.9

1. $2\tanh x\,\text{sech}^2 x$ 3. $\text{sech}\,x\,\text{csch}\,x$ 5. $\frac{2e^{2x}\cosh(\arctan e^{2x})}{1+e^{4x}}$ 7. $\frac{-x\,\text{csch}^2\sqrt{x^2+1}}{\sqrt{x^2+1}}$ 9. $\frac{-\text{sech}\,x(\tanh x + 2\sinh x)}{(1+\cosh x)^2}$

15. (a) $\frac{3}{5}$ (b) $\frac{4}{5}$ (c) $\frac{4}{3}$ (d) $\frac{5}{4}$ (e) $\frac{1}{4}$

25. (a) absolute max $f(0) = 1$
(b) points of inflection at $x = \ln(1+\sqrt{2}) \cong 0.881$, $x = -\ln(1+\sqrt{2}) \cong -0.881$
(c) concave up on $(-\infty, -\ln(1+\sqrt{2})) \cup (\ln(1+\sqrt{2}), \infty)$; concave down on $(-\ln(1+\sqrt{2}), \ln(1+\sqrt{2}))$

(d)

Download full file from answersun.com