

Student Solutions Manual
for use with
**Complex Variables
and Applications**
Seventh Edition

Selected Solutions to Exercises in Chapters 1-7

by

James Ward Brown
Professor of Mathematics
The University of Michigan- Dearborn

Ruel V. Churchill
Late Professor of Mathematics
The University of Michigan



Higher Education

Boston Burr Ridge, IL Dubuque, IA Madison, WI New York San Francisco St. Louis
Bangkok Bogotá Caracas Kuala Lumpur Lisbon London Madrid Mexico City
Milan Montreal New Delhi Santiago Seoul Singapore Sydney Taipei Toronto

Table of Contents

Chapter 1 1

Chapter 2 22

Chapter 3 35


Chapter 4 53

Chapter 5 75


Chapter 6 94

Chapter 7 118

(c) $z_1 = (-3, 1), \quad z_2 = (1, 4)$



(d) $z_1 = x_1 + iy_1, \quad z_2 = x_1 - iy_1$



2. Inequalities (3), Sec. 4, are

$$\operatorname{Re} z \leq |\operatorname{Re} z| \leq |z| \quad \text{and} \quad \operatorname{Im} z \leq |\operatorname{Im} z| \leq |z|.$$

These are obvious if we write them as

$$x \leq |x| \leq \sqrt{x^2 + y^2} \quad \text{and} \quad y \leq |y| \leq \sqrt{x^2 + y^2}.$$

3. In order to verify the inequality $\sqrt{2}|z| \geq |\operatorname{Re} z| + |\operatorname{Im} z|$, we rewrite it in the following ways:

$$\sqrt{2}\sqrt{x^2 + y^2} \geq |x| + |y|,$$

$$2(x^2 + y^2) \geq |x|^2 + 2|x||y| + |y|^2,$$

$$|x|^2 - 2|x||y| + |y|^2 \geq 0,$$

$$(|x| - |y|)^2 \geq 0.$$

This last form of the inequality to be verified is obviously true since the left-hand side is a perfect square.

6. The four roots of the equation $z^4 + 4 = 0$ are the four fourth roots of the number -4 . To find those roots, we write $-4 = 4 \exp[i(\pi + 2k\pi)]$ ($k = 0, \pm 1, \pm 2, \dots$). Then

$$(-4)^{1/4} = \sqrt{2} \exp\left[i\left(\frac{\pi}{4} + \frac{k\pi}{2}\right)\right] = \sqrt{2} e^{i\pi/4} e^{ik\pi/2} \quad (k = 0, 1, 2, 3).$$

To be specific,

$$c_0 = \sqrt{2} e^{i\pi/4} = \sqrt{2} \left(\cos \frac{\pi}{4} + i \sin \frac{\pi}{4} \right) = \sqrt{2} \left(\frac{1}{\sqrt{2}} + i \frac{1}{\sqrt{2}} \right) = 1 + i,$$

$$c_1 = c_0 e^{i\pi/2} = (1 + i)i = -1 + i,$$

$$c_2 = c_0 e^{i\pi} = (1 + i)(-1) = -1 - i,$$

$$c_3 = c_0 e^{i3\pi/2} = (1 + i)(-i) = 1 - i.$$

This enables us to write

$$\begin{aligned} z^4 + 4 &= (z - c_0)(z - c_1)(z - c_2)(z - c_3) \\ &= [(z - c_1)(z - c_2)] \cdot [(z - c_0)(z - c_3)] \\ &= [(z + 1) - i][(z + 1) + i] \cdot [(z - 1) - i][(z - 1) + i] \\ &= [(z + 1)^2 + 1] \cdot [(z - 1)^2 + 1] \\ &= (z^2 + 2z + 2)(z^2 - 2z + 2). \end{aligned}$$

7. Let c be any n th root of unity other than unity itself. With the aid of the identity (see Exercise 10, Sec. 7),

$$1 + z + z^2 + \dots + z^{n-1} = \frac{1 - z^n}{1 - z} \quad (z \neq 1),$$

we find that

$$1 + c + c^2 + \dots + c^{n-1} = \frac{1 - c^n}{1 - c} = \frac{1 - 1}{1 - c} = 0.$$

9. Observe first that

$$(z^{1/m})^{-1} = \left[\sqrt[m]{r} \exp \frac{i(\theta + 2k\pi)}{m} \right]^{-1} = \frac{1}{\sqrt[m]{r}} \exp \frac{i(-\theta - 2k\pi)}{m} = \frac{1}{\sqrt[m]{r}} \exp \frac{i(-\theta)}{m} \exp \frac{i(-2k\pi)}{m}$$

If the Cauchy-Riemann equations $u_x = v_y$, $u_y = -v_x$ hold, then

$$\sin x \cosh y = 0 \quad \text{and} \quad \cos x \sinh y = 0.$$

The first of these equations tells us that $\sin x = 0$, or $x = n\pi$ ($n = 0, \pm 1, \pm 2, \dots$). Since $\cos n\pi \neq 0$, it follows that $\sinh y = 0$, or $y = 0$. Consequently, the Cauchy-Riemann equations hold only when

$$z = n\pi \quad (n = 0, \pm 1, \pm 2, \dots).$$

So there is no neighborhood throughout which f is analytic, and this means that $\cos \bar{z}$ is nowhere analytic.

16. (a) Use expression (12), Sec. 33, to write

$$\overline{\cos(iz)} = \overline{\cos(-y + ix)} = \cos y \cosh x - i \sin y \sinh x$$

and

$$\cos(i\bar{z}) = \cos(y + ix) = \cos y \cosh x - i \sin y \sinh x.$$

This shows that $\overline{\cos(iz)} = \cos(i\bar{z})$ for all z .

(b) Use expression (11), Sec. 33, to write

$$\overline{\sin(iz)} = \overline{\sin(-y + ix)} = -\sin y \cosh x - i \cos y \sinh x$$

and

$$\sin(i\bar{z}) = \sin(y + ix) = \sin y \cosh x + i \cos y \sinh x.$$

Evidently, then, the equation $\overline{\sin(iz)} = \sin(i\bar{z})$ is equivalent to the pair of equations

$$\sin y \cosh x = 0, \quad \cos y \sinh x = 0.$$

Since $\cosh x$ is never zero, the first of these equations tells us that $\sin y = 0$. Consequently, $y = n\pi$ ($n = 0, \pm 1, \pm 2, \dots$). Since $\cos n\pi = (-1)^n \neq 0$, the second equation tells us that $\sinh x = 0$, or that $x = 0$. So we may conclude that $\overline{\sin(iz)} = \sin(i\bar{z})$ if and only if $z = 0 + in\pi = n\pi i$ ($n = 0, \pm 1, \pm 2, \dots$).


17. Rewriting the equation $\sin z = \cosh 4$ as $\sin x \cosh y + i \cos x \sinh y = \cosh 4$, we see that we need to solve the pair of equations

$$\sin x \cosh y = \cosh 4, \quad \cos x \sinh y = 0$$

(b) Here $C: z = x$ ($0 \leq x \leq 2$). Then

$$\int_C (z-1) dz = \int_0^2 (x-1) dx = \left[\frac{x^2}{2} - x \right]_0^2 = 0.$$

3. In this problem, the path C is the sum of the paths C_1 , C_2 , C_3 , and C_4 that are shown below.



The function to be integrated around the closed path C is $f(z) = \pi e^{\pi \bar{z}}$. We observe that $C = C_1 + C_2 + C_3 + C_4$ and find the values of the integrals along the individual legs of the square C .

(i) Since C_1 is $z = x$ ($0 \leq x \leq 1$),

$$\int_{C_1} \pi e^{\pi \bar{z}} dz = \pi \int_0^1 e^{\pi x} dx = e^{\pi} - 1.$$

(ii) Since C_2 is $z = 1 + iy$ ($0 \leq y \leq 1$),

$$\int_{C_2} \pi e^{\pi \bar{z}} dz = \pi \int_0^1 e^{\pi(1-iy)} i dy = e^{\pi} \pi i \int_0^1 e^{-i\pi y} dy = 2e^{\pi}.$$

(iii) Since C_3 is $z = (1-x) + i$ ($0 \leq x \leq 1$),

$$\int_{C_3} \pi e^{\pi \bar{z}} dz = \pi \int_0^1 e^{\pi[(1-x)-i]} (-1) dx = \pi e^{\pi} \int_0^1 e^{-\pi x} dx = e^{\pi} - 1.$$

(iv) Since C_4 is $z = i(1-y)$ ($0 \leq y \leq 1$),

$$\int_{C_4} \pi e^{\pi \bar{z}} dz = \pi \int_0^1 e^{-\pi(1-y)i} (-i) dy = \pi i \int_0^1 e^{i\pi y} dy = -2.$$

Finally, then, since

$$\int_C \pi e^{\pi \bar{z}} dz = \int_{C_1} \pi e^{\pi \bar{z}} dz + \int_{C_2} \pi e^{\pi \bar{z}} dz + \int_{C_3} \pi e^{\pi \bar{z}} dz + \int_{C_4} \pi e^{\pi \bar{z}} dz,$$

we find that

$$\int_C \pi e^{\pi \bar{z}} dz = 4(e^{\pi} - 1).$$

78

2. (b) Replacing z by $z - 1$ in the known expansion

$$e^z = \sum_{n=0}^{\infty} \frac{z^n}{n!} \quad (|z| < \infty),$$

we have

$$e^{z-1} = \sum_{n=0}^{\infty} \frac{(z-1)^n}{n!} \quad (|z| < \infty).$$

So

$$e^z = e^{z-1} e = e \sum_{n=0}^{\infty} \frac{(z-1)^n}{n!} \quad (|z| < \infty).$$

3. We want to find the Maclaurin series for the function

$$f(z) = \frac{z}{z^4 + 9} = \frac{z}{9} \cdot \frac{1}{1 + (z^4/9)}.$$

To do this, we first replace z by $-(z^4/9)$ in the known expansion

$$\frac{1}{1-z} = \sum_{n=0}^{\infty} z^n \quad (|z| < 1),$$

as well as its condition of validity, to get

$$\frac{1}{1 + (z^4/9)} = \sum_{n=0}^{\infty} \frac{(-1)^n}{3^{2n}} z^{4n} \quad (|z| < \sqrt{3}).$$

Then, if we multiply through this last equation by $\frac{z}{9}$, we have the desired expansion:

$$f(z) = \sum_{n=0}^{\infty} \frac{(-1)^n}{3^{2n+2}} z^{4n+1} \quad (|z| < \sqrt{3}).$$

6. Replacing z by z^2 in the representation

$$\sin z = \sum_{n=0}^{\infty} (-1)^n \frac{z^{2n+1}}{(2n+1)!} \quad (|z| < \infty),$$

we have

$$\sin(z^2) = \sum_{n=0}^{\infty} (-1)^n \frac{z^{4n+2}}{(2n+1)!} \quad (|z| < \infty).$$

(b) When $f(z) = \frac{1}{1+z^2}$, we have

$$\frac{1}{z^2} f\left(\frac{1}{z}\right) = \frac{1}{1+z^2} = \frac{1}{1-(-z^2)} = 1 - z^2 + z^4 - \dots \quad (0 < |z| < 1).$$

Thus

$$\int_C f(z) dz = 2\pi i \operatorname{Res}_{z=0} \frac{1}{z^2} f\left(\frac{1}{z}\right) = 2\pi i(0) = 0.$$

(c) If $f(z) = \frac{1}{z}$, it follows that $\frac{1}{z^2} f\left(\frac{1}{z}\right) = \frac{1}{z}$. Evidently, then,

$$\int_C f(z) dz = 2\pi i \operatorname{Res}_{z=0} \frac{1}{z^2} f\left(\frac{1}{z}\right) = 2\pi i(1) = 2\pi i.$$

4. Let C denote the circle $|z|=1$, taken counterclockwise.

(a) The Maclaurin series $e^z = \sum_{n=0}^{\infty} \frac{z^n}{n!}$ ($|z| < \infty$) enables us to write

$$\int_C \exp\left(z + \frac{1}{z}\right) dz = \int_C e^z e^{1/z} dz = \int_C e^{1/z} \sum_{n=0}^{\infty} \frac{z^n}{n!} dz = \sum_{n=0}^{\infty} \frac{1}{n!} \int_C z^n \exp\left(\frac{1}{z}\right) dz.$$

(b) Referring to the Maclaurin series for e^z once again, let us write

$$z^n \exp\left(\frac{1}{z}\right) = z^n \sum_{k=0}^{\infty} \frac{1}{k!} \cdot \frac{1}{z^k} = \sum_{k=0}^{\infty} \frac{1}{k!} z^{n-k} \quad (n = 0, 1, 2, \dots).$$

Now the $\frac{1}{z}$ in this series occurs when $n-k=-1$, or $k=n+1$. So, by the residue theorem,

$$\int_C z^n \exp\left(\frac{1}{z}\right) dz = 2\pi i \frac{1}{(n+1)!} \quad (n = 0, 1, 2, \dots).$$

The final result in part (a) thus reduces to


$$\int_C \exp\left(z + \frac{1}{z}\right) dz = 2\pi i \sum_{n=0}^{\infty} \frac{1}{n!(n+1)!}.$$

6. (a) Let us first use the branch

$$f(z) = \frac{z^{-1/2}}{z^2 + 1} = \frac{\exp\left(-\frac{1}{2}\log z\right)}{z^2 + 1} \quad \left(|z| > 0, -\frac{\pi}{2} < \arg z < \frac{3\pi}{2}\right)$$

and the indented path shown below to evaluate the improper integral

$$\int_0^{\infty} \frac{dx}{\sqrt{x}(x^2 + 1)}.$$



Cauchy's residue theorem tells us that

$$\int_{L_1} f(z) dz + \int_{C_R} f(z) dz + \int_{L_2} f(z) dz + \int_{C_\rho} f(z) dz = 2\pi i \operatorname{Res}_{z=i} f(z),$$

or

$$\int_{L_1} f(z) dz + \int_{L_2} f(z) dz = 2\pi i \operatorname{Res}_{z=i} f(z) - \int_{C_\rho} f(z) dz - \int_{C_R} f(z) dz.$$

Since

$$L_1: z = re^{i0} = r \quad (\rho \leq r \leq R) \quad \text{and} \quad -L_2: z = re^{i\pi} = -r \quad (\rho \leq r \leq R),$$

we may write

$$\int_{L_1} f(z) dz + \int_{L_2} f(z) dz = \int_{\rho}^R \frac{dr}{\sqrt{r}(r^2 + 1)} - i \int_{\rho}^R \frac{dr}{\sqrt{r}(r^2 + 1)} = (1 - i) \int_{\rho}^R \frac{dr}{\sqrt{r}(r^2 + 1)}.$$

Thus

$$(1 - i) \int_{\rho}^R \frac{dr}{\sqrt{r}(r^2 + 1)} = 2\pi i \operatorname{Res}_{z=i} f(z) - \int_{C_\rho} f(z) dz - \int_{C_R} f(z) dz.$$